

Autoimmune/inflammatory syndrome induced by adjuvants (ASIA) 2013: Unveiling the pathogenic, clinical and diagnostic aspects

Carlo Perricone ¹, Serena Colafrancesco, Roei D Mazor, Alessandra Soriano, Nancy Agmon-Levin, Yehuda Shoenfeld

Affiliations

PMID: 24238833 DOI: 10.1016/j.jaut.2013.10.004

Abstract

In 2011 a new syndrome termed 'ASIA Autoimmune/Inflammatory Syndrome Induced by Adjuvants' was defined pointing to summarize for the first time the spectrum of immune-mediated diseases triggered by an adjuvant stimulus such as chronic exposure to silicone, tetramethylpentadecane, pristane, aluminum and other adjuvants, as well as infectious components, that also may have an adjuvant effect. All these environmental factors have been found to induce autoimmunity by themselves both in animal models and in humans; for instance, silicone was associated with siliconosis, aluminum hydroxide with post-vaccination phenomena and macrophagic myofasciitis syndrome. Several mechanisms have been hypothesized to be involved in the onset of adjuvant-induced autoimmunity; a genetic favorable background plays a key role in the appearance on such vaccine-related diseases and also justifies the rarity of these phenomena. This paper will focus on protean facets which are part of ASIA, focusing on the roles and mechanisms of action of different adjuvants which lead to the autoimmune/inflammatory response. The data herein illustrate the critical role of environmental factors in the induction of autoimmunity. Indeed, it is the interplay of genetic susceptibility and environment that is the major player for the initiation of breach of tolerance.

Keywords: Adjuvant; Autoantibodies; Autoimmune/Inflammatory syndrome induced by adjuvants; Autoimmunity; *Saccharomyces cerevisiae*; Vaccine.

Crown Copyright © 2013. Published by Elsevier Ltd. All rights reserved.